


# Better Specimen Collection Better Diagnostics





COPAN FLOQSwabs<sup>®</sup> consist of molded plastic applicator shafts with tips (in varying dimensions) coated with Nylon<sup>®</sup> fibers, creating a thin absorbent layer allowing for quick sample uptake and elution of more than 90% of the sample.<sup>1</sup> The unique swab design ensures the most effective specimen collection to expand downstream testing capabilities.<sup>2</sup>



### Superior sample collection

FLOQSwabs<sup>®</sup> have no internal core to trap the sample, allowing rapid absorption and ensuring fast and complete specimen elution.

### **Designed for multiple** collection sites

Various shaft and tip dimensions make FLOQSwabs<sup>®</sup> an improved alternative to invasive and costly collection procedures.<sup>4, 5</sup>

## The perfect partner

FLOQSwabs<sup>®</sup> are compatible with a variety of bacteriology and virology transport media, ensuring broad assay compatibility.<sup>2, 3, 6-10</sup>

# Ϋ́Ω Why Flocked?

Maximizing sample recovery allows multiple tests to be run from a single collection, and improves the sample yield from each collection, improving efficiency and diagnostic accuracy.

# Traditional Fiber Swab

Yards of fiber wrapped around applicator



#### Patent Information

US 8,317,728 8,114,027 8,979,784 9,011,358 13/657,949 16/448,398 9,173,779 14/448398 10,327,741 | EP1608268 | JP4579902 | AU2004226798 | CA2,515,205 | NZ541560 | SG170654 | CNZL200610099310.9

# **FLOQSwabs**®

Nylon<sup>®</sup> fibers are applied to applicator using a proprietary flocking process

> More of sample available for testing

Sample diffuses and becomes trapped in the fibers.

Sample stays close to the surface, eluting upon contact with transport media. Brush like texture dislodges and collects cells.

How many tests from one sample?

Multiple aliquots for various tests<sup>6-11</sup>

# Ordering Information

FLOQSwabs<sup>®</sup> are available with various tip sizes and breakpoints to be used for various applications.

All FLOQSwabs<sup>®</sup> are packaged sterile and come in boxes of 100 pieces, 10 boxes to a case, unless otherwise noted.

Learn More

## FLOQSwabs® in Peel Pouch

FLEXIBLE					
	534CS01	Commonly Used For	Break Point	Product	Cat. No.
MINI FLEXIB	516CS01	Nasal, throat, vaginal, groin, axilla, rectal, wound, buccal, stool	100 mm	REGULAR	519C501
LONG CONTOURED WI	56380CS01	Nasal, throat, vaginal, groin, axilla, rectal, wound, buccal, stool	80 mm	REGULAR	502CS01
CONTOURED WITH LO	56780CS01	Nasal, throat, groin, axilla, wound, buccal, stool	30 mm	REGULAR	520CS01
SHORT CONTOURED W	56750CS01	Urethral, nasal, pediatric	100 mm	MINITIP	518CS01
		Urethral, nasal, pediatric	80 mm	MINITIP	501CS01

# FLOQSwabs® in Peel Pouch

Cat.	No.	Product
5250	501	ULTRA-THIN MINITIP
503C	501	FLEXIBLE
534C	501	FLEXIBLE
516CS01	501	MINI FLEXIBLE
		LONG CONTOURED WITH ST
56380CS01	501	
56700	5501	CONTOURED WITH LOW STO
56780CS01		
		SHORT CONTOURED WITH S
56750CS01	CS01	

	Break Point	Commonly Used For
TIP	80 mm	Urethral
	100 mm	Nasopharyngeal
	N/A	Nasopharyngeal
	100 mm	Neonatal
H STOPPER	80 mm	Mid Turbinate
STOPPER	80 mm	Mid Turbinate
H STOPPER	50 mm	Mid Turbinate

#### Break Commonly Cat. No. **Product** Used For Point REGULAR Nasal, throat, vaginal, 519C 100 mm groin, axilla, rectal, COPAN €€ 0123 § FLOQSwabs\* wound, buccal, stool REGULAR Nasal, throat, vaginal, 552C 80 mm groin, axilla, rectal, COPAN FLOOSwabs" CE 0123 6 wound, buccal, stool REGULAR Nasal, throat, groin, 520C 30 mm FLOQSwabs\* COPAN C€ 0123 § axilla, buccal, stool **REGULAR WITH LOW STOPPER** 5U002S 80 mm Rectal, entrics COPAN C€ 0123 § FLOOSwabs' LARGE TIP WITH STOPPER 528C N/A Buccal FLOQSwabs' COPAN CE 0123 5 MINITIP Urethral, nasal, 518C 100 mm pediatric COPAN (€ 0123 § FLOQSwabs" MINITIP Urethral, nasal, 551C 80 mm pediatric FLOQSwabs COPAN C€ 0123 §

FLOQSwabs<sup>®</sup> in Tube

### FLOQSwabs<sup>®</sup> in Tube



#### **Scientific References**

- 1. Copan Group. Data on file Internal elution studies.
- З. Copan Italia S.p.A. (2020). FLOQSwabs® in pouch: Product insert & how to use guide (HPC217 Rev.00). Copan Italia S.p.A.
- Saville, M., Hawkes, D., et al. (2020). Analytical performance of HPV assays on vaginal self-collected vs practitioner-collected cervical samples: The 4. SCoPE study. Journal of Clinical Virology, 127, 104375. https://doi.org/10.1016/j.jcv.2020.104375
- 5. Misra, A., Speicher, D. J., et al. (2021). Self-collected oral flocked swabs to measure prevalence of Epstein-Barr Virus antibodies and DNA amongst university students. Diagnostic Microbiology and Infectious Disease, 100, 115295. https://doi.org/10.1016/j.diagmicrobio.2020.115295 Van, T. T., Cox, L. M., et al. (2017). Prevalence of Fusobacterium necrophorum in children presenting with pharyngitis. Journal of Clinical Microbiology,
- 6. 55(4), 1147–1153. https://doi.org/10.1128/JCM.02174-16
- Cleveland Clinic Laboratories. (2024). Microbiology specimen collection & transport guidelines. Cleveland Clinic. Retrieved from https:// 7. clevelandcliniclabs.com/laboratory-resources/specimen-collection-transport/
- Fontana, C., Favaro, M., & Favalli, C. (2013). How liquid-based microbiology can change the workflow in microbiology laboratories. Advances in 8 Microbiology, 3(6), 504-510. https://doi.org/10.4236/aim.2013.36067
- 9. Plapp, F. (2014). E Swab Collection System. ClinLab Navigator. https://www.clinlabnavigator.com/blog/e-swab-collection-system 10. Chernesky, M., Castriciano, S., et al. (2006). Use of flocked swabs and a universal transport medium to enhance molecular detection of Chlamydia
- 11. Copan Diagnostics Inc. (2007). ESwab: New liquid-based transport system for microbiology with flocked swabs. Retrieved from Rapid Microbiology website: https://www.rapidmicrobiology.com

	Break Point	Commonly Used For
	100 mm	Nasopharyngeal
CC 0123 S	100 mm	Neonatal
	80 mm	Urethral

Munywoki, P. K., Hamid, F. et al. (2011). Improved detection of respiratory viruses in pediatric outpatients with acute respiratory illness by real-time PCR using nasopharyngeal flocked swabs. Journal of Clinical Microbiology, 49(9), 3365–3367. https://doi.org/10.1128/JCM.02231-10

trachomatis and Neisseria gonorrhoeae. Journal of Clinical Microbiology, 44(3), 1084–1086. https://doi.org/10.1128/JCM.44.3.1084-1086.2006



These products may not be available in all Countries. Please refer to www.copanusa.com for the most recent version of this document. Always consult product technical documentation for complete information. The use of this product in association with diagnostic kits or instrumentation should be validated by the end user. All trademarks, logos and brand names are the property of their respective owners. ©2025 COPAN Diagnostics Inc. All rights reserved. Code: FLOQ Brochure\_03\_2025



**COPAN Diagnostics, Inc.** 26055 Jefferson Avenue | Murrieta, CA 92562 USA